75 per cent in Shuqra. In most localities, the infection rates ranged from 27 to 88 per cent among children under 15 years of age and from zero to 48 per cent in persons over 15 years of age. The Fadhli and the Abdali tribal districts were the most seriously affected regions visited.

Diseases of the Skin. Favus, ringworm and other mycotic skin infections are widely distributed. Madura foot is common in the Protectorate, and cases are seen in the Colony hospitals each year among patients from the interior. Phagedenic ulcers are responsible for considerable disability among laborers in the Colony and in all parts of the Protectorate. Several thousand cases are treated each year in the hospitals and the dispensaries in Aden Colony, where they are usually designated as "Yemen ulcers." The ankle is the most frequent site of infection, and amputation is often necessary in neglected cases.

Scabies and impetigo are prevalent, especially among the children of inland towns and villages who live under congested, insanitary conditions. By contrast, surprisingly few cases of scabies are encountered among the children seen in the schools in Aden township.

Leprosy. Leprosy is endemic in the Protectorate, but no indigenous cases have been reported from the Colony within recent years. The infection is probably not extensive, although foci of the disease are known to exist in the Hadhramaut and in the Lahej, the Dhala' and the Abyen areas in the western protectorate. As lepers are not discriminated against socially, some difficulty is encountered in controlling the movement of infected individuals in the Colony. A small leper hospital is located at Sheikh Othman: it receives patients not only from Aden territory but also from Yemen. Eighteen of the 29 patients in residence in 1939[18] were open nodular or lepromatous cases.

Other Infections. Tetanus is endemic, but reported cases are rare.

p. 249(A)